

ISSN 0556-3321
CODEN PJB 0B6

PAKISTAN JOURNAL OF BOTANY

A BIMONTHLY PUBLICATION OF PAKISTAN BOTANICAL SOCIETY

Volume 38 No. 5 (Special Issue), December, 2006

Guest Editors

Muhammad Ashraf

Department of Botany, University of Agriculture, Faisalabad, Pakistan
And

Habib-ur-Rehman Athar

Institute of Pure and Applied Biology, Bahauddin Zakariya University, Multan, Pakistan

Chief Editor:

Dr. Abdul Ghaffar

Department of Botany, University of Karachi, Karachi-75270, Pakistan.

E-mail: shaji@super.net.pk

Subject Editors:

Plant Biosystematics, Taxonomy and Evolution -Dr. M. Qaiser

Department of Botany, University of Karachi, Karachi-75270, Pakistan.

E-mail: flora@super.net.pk

Plant Biochemistry and Physiology -Dr. M. Ashraf

Department of Botany, University of Agriculture, Faisalabad, Pakistan.

E-mail: ashrafbot@yahoo.com

Ecology, Phycology and Biodiversity -Dr. Farrukh Hussain

Department of Botany, University of Peshawar, Peshawar, Pakistan.

E-mail: Farrukhbotany@yahoo.com

Cell Biology, Genetics and Microbiology -Dr. Shahida Hasnain

Department of Botany, University of Punjab, Lahore, Pakistan.

E-mail: genetic@brain.net.pk

Mycology and Plant Pathology -Dr. Abdul Ghaffar

Department of Botany, University of Karachi, Karachi-75270, Pakistan.

E-mail: shaji@super.net.pk

Editorial Committee: Dr. S.I. Ali, Dr. Mustafa Shameel, Dr. S. Shahid Shaukat (University of Karachi), Dr. Iftikhar Ahmed (PARC, Islamabad), Dr. Kausar A. Malik (PAEC, Islamabad), Dr. Javed Iqbal (Punjab University, Lahore), Dr. Riaz H. Qureshi (University of Agric., Faisalabad), Mr. I.C. Hedge (Royal Botanical Garden, Edinburgh, UK), Dr. Klaus Lüning (Stiftung Alfred-Werger-Institut, Germany), Dr. Irwin A. Ungar (Ohio University, USA), Dr. M.A. El-Wakil (University of Mansoura, Egypt)

COMPARATIVE EFFECTS OF NaCl AND SEASALT ON SEED GERMINATION OF COASTAL HALOPHYTES

ABDUL HAMEED, MUHAMMAD ZAHEER AHMED AND MUHAMMAD AJMAL KHAN*

Department of Botany, University of Karachi, Karachi-75270, Pakistan

Abstract

The coastal and near coastal habitats of Karachi, Pakistan are characterized by dominant stands of perennial halophytes like *Aeluropus lagopoides*, *Desmostachya bipinnata*, *Haloxylon stocksii* and *Suaeda fruticosa*. Experiments were carried out to investigate comparative effects of NaCl and seasalt salinity on both rate and final seed germination of these species. Salinity tolerance range of halophytes varied among species and also between salt types. Seed germination of all test species decreased with increase in salinity of both salt types. *Aeluropus lagopoides* was the most tolerant at 50 dS m⁻¹ whereas, *S. fruticosa* the least salt tolerant at 30 dS m⁻¹, although the tolerance range was in following sequence: *A. lagopoides* > *D. bipinnata* > *H. stocksii* > *S. fruticosa*. While rate of germination showed a different pattern *D. bipinnata* > *A. lagopoides* = *H. stocksii* > *S. fruticosa*. Seasalt affected seed germination as well as rate of germination of *A. lagopoides*, *D. bipinnata* and *S. fruticosa* more adversely than NaCl especially at higher concentration. However, in *H. stocksii* seed germination was not significantly different in either salt type. Rate of germination also followed same pattern

Keywords: *Aeluropus lagopoides*, *Desmostachya bipinnata*, *Haloxylon stocksii*, Karachi, *Suaeda fruticosa*

Introduction

Halophytes are distributed in coastal and inland saline habitats and face direct and secondary effects of salinity stress at every stage of their life cycles (Adam, 1990; Khan & Gul, 2002a; Khan & Qaiser, 2006; Ungar, 1991). Germination is reported to be more vulnerable stage to salinity than others (Khan & Gul, 2002a, 2006). Halophytes show optimal seed germination under non-saline conditions like glycophytes but differ in their ability to germinate in presence of even higher levels of salinity (Ungar, 1995; Noe & Zedler, 2000; Khan & Gul, 2002a). Salinity tolerance of halophytes at germination varies among species (Yokoishi & Tanimoto, 1994; Ungar, 1995; Khan *et al.* 2002, 2006; Liu *et al.* 2006). Threshold tolerance level to salt stress at germination stage depends on the type of species, e.g., for *Salicornia herbacea* it is 1700 mM (Chapman, 1960) while *Spartina alterniflora* can tolerate 1030 mM NaCl stress (Mooring *et al.*, 1971). Khan & Gul (2006) reviewed that approximately 83 % Dicotyledonous halophytes can germinate at or above sea water salinity (60 dS m⁻¹), whereas only 17 % of monocotyledonous species can tolerate such a high salinity at germination stage. However, information about seed germination of halophytes is far from complete, since actual halophytic diversity is unknown (Khan & Qaiser, 2006). Many of these can be potential candidates for producing food, fodder or forage using saline agriculture, which is likely to stay in future. However, success of saline agriculture greatly depends upon the knowledge of seed

* Corresponding author: Muhammad Ajmal Khan; halophyte_ajmal@yahoo.com

germination eco-physiology of halophytes, since most of halophytic species propagate through seeds.

Most of the studies on effect of salinity on seed germination of halophytes have been conducted using NaCl solutions. Such investigations may not infer germination responses of plants under field conditions, because in field soil contains different salts, which collectively influence seed germination in a different way from their individual effects (Ungar, 1978). Seasalt however mimic in composition to soil solution and can be used to analyze synergistic effect of different salts on seed germination (Liu *et al.* 2006). Unfortunately, little is known about how seawater affects on seed germination of halophytes (Baskin & Baskin, 1998; Houle *et al.* 2001; Khan *et al.* 2006). It appears that seawater inhibits germination more in comparison to NaCl (De Villiers *et al.* 1994; Houle *et al.* 2001; Zia & Khan, 2002). However, Tirmizi *et al.* (1993) found that NaCl inhibited the germination of *Hipophae rhamnoides* more than seawater. Similarly Atia *et al.* (2006) also reported that NaCl inhibits seed germination more in comparison to sea-salt.

Pakistani coastal areas are inhabited by 108 halophytic species belonging to 36 families. Most are perennial including grasses (20), sedges (14) and chenopods (13) (Khan & Gul, 2002b). *Aeluropus lagopoides*, *Desmostachya bipinnata*, *Haloxylon stocksii*, and *Suaeda fruticosa* are halophytes found in coastal habitats of Karachi, Pakistan. These halophytic species with wide ecological range are of medicinally important or used as forage or could also be used as a coastal dune stabilizer. Study under consideration was designed to investigate comparative effects of NaCl and seasalt salinity on seed germination of these potentially important halophytes.

Material and Methods

Seeds of test species were collected from a various saline habitats of Karachi Coast during March to April, 2005. Seeds were separated from the inflorescence and dry - stored at room temperature. Before storage they were surface sterilized using 0.85 % Clorox solution for one minute followed by thorough rinsing with autoclaved distilled water and air-drying. Germination was carried out in 5 cm diameter, tight-fitting plastic Petri dishes with 5 ml of test solution. Germination was carried out in NaCl and seawater (0, 10, 20, 30 40 and 50 dS m⁻¹) separately. Four replicates of 25 seeds each were used for each treatment. Seeds were considered to be germinated with radicle emergence. Seeds were germinated in an incubator at a day/night temperature of 20/30 °C with 12-h photoperiod (Sylvania cool white fluorescent lamps, 25 μmol m⁻² s⁻¹, 400-750 nm). Percent germination was recorded on alternate days for 20 days. The rate of germination was estimated by using modified Timson's index of germination velocity which is as follows: $\Sigma G/t$; where G is percentage of seed germination at 2-d intervals, and t is total germination period (Khan & Ungar, 1984). The maximum value possible for our data using this index was 50 (i.e. 1000/20). The higher the value more rapid the germination. Germination data was arcsine transformed before statistical analysis to ensure homogeneity of variance. Entire experiment was repeated twice. These data were analyzed using SPSS for Windows release 11 (SPSS 2001). The effect of NaCl and seasalt on the germination and rate of germination of test species was examined using analysis of variance (ANOVA). A Bonferonni post hoc test was used to determine significant differences ($p < 0.05$) between means.

Results

Three-way analyses of variance show that the interaction of species (SP), salt type (ST) and concentration (CON) were significantly differed, although their individual effects except concentration were non-significant for both final ($F = 1.97$, $P < 0.01$) and rate ($F = 1.80$, $P < 0.01$) of seed germination. The interactive effect of SP and ST was also significant at both final ($F = 9.17$, $P < 0.001$) and rate ($F = 8.40$, $P < 0.001$) of seed germination level (Table 1).

In non-saline control *Aeluropus lagopoides* and *Desmostachya bipinnata* showed 100 % seed germination whereas only 80 % seeds germinated for *Haloxylon stocksii* and *Suaeda fruticosa* (Figure 1). Increase in salinity linearly decreased germination for *H. stocksii* and *S. fruticosa*, while in *A. lagopoides* and *D. bipinnata* threshold salinity for germination suppression was 20 dS m^{-1} . Salinity tolerance limit of test species decreased in following sequence: *A. lagopoides* > *D. bipinnata* > *H. stocksii* > *S. fruticosa*.

Table 1. Three-way ANOVA showing significance of the effect of various factors on final and rate of germination of halophytes.

	Source	Type III Sum of squares	df	F
Final Germination	SPECIES (SP)	25527	3	4.32 ^{ns}
	SALT TYPE (ST)	5188	1	1.81 ^{ns}
	CONCENTRATION (CON)	157678	5	22.23 ^{**}
	SP × ST	5424	3	9.17 ^{**}
	SP × CON	5361	15	1.81 ^{ns}
	ST × CON	6294	5	6.40 ^{**}
	SP × ST × CON	2956	15	1.97 [*]
Rate of Germination	SPECIES (SP)	2098	3	3.25 ^{ns}
	SALT TYPE (ST)	428	1	1.55 ^{ns}
	CONCENTRATION (CON)	31354	5	50.33 ^{***}
	SP × ST	588	3	8.40 ^{**}
	SP × CON	644	15	1.83 ^{ns}
	ST × CON	525	5	4.50 [*]
	SP × ST × CON	351	15	1.80 [*]

Where, ns = non-significant, * = $P < 0.01$, ** = $P < 0.001$ and *** = $P < 0.0001$.




Fig. 1. Effect of NaCl and Seasalt on mean final seed germination of coastal halophytes.




Fig. 2. Effect of NaCl and Seasalt on rate of seed germination of coastal halophytes.

For *A. lagopoides* mean final germination was not significantly different ($P < 0.05$) from non-saline control in either salt type up to 20 dS m^{-1} , but a further increase in salinity linearly decreased germination in both salts. However, seasalt decreased germination more in comparison to NaCl and 25% and 10 % seeds germinated at 50 dS m^{-1} of NaCl and seasalt respectively (Figure 1a). Seed germination of *D. bipinnata* linearly decreased with increasing seasalt salinity, but it was non-significantly ($P < 0.05$) different from control in NaCl up to 30 dS m^{-1} and at 50 dS m^{-1} of seasalt there was no seed germination, while 20 % seeds germinated at similar salinity level of NaCl (Figure 1b). In *H. stocksii* mean final germination progressively decreased with increase in salinity, with no germination above 50 dS m^{-1} and it was also not affected significantly ($P < 0.05$) by salt type (Figure 1c). However, mean final germination in *S. fruticosa* significantly ($P < 0.05$) differed between salt types, only above 10 dS m^{-1} with comparatively more adverse effects in seasalt than NaCl and no seed germination was found above 30 dS m^{-1} of both salts (Figure 1d).

Rate of seed germination for all test species was highest in distilled water controls. Increasing salinity linearly decreased rate of germination, however, seasalt delayed rate of germination more than NaCl at or above 30 dS m^{-1} in all except *H. stocksii*. Rate of germination of test species was in following order: *D. bipinnata* > *A. lagopoides* = *H. stocksii* > *S. fruticosa* (Figure 2).

Discussion

Salinity tolerance range of halophytes varied among species. Khan & Gul (2002) categorized the halophytes and found that halophytic species around the Karachi coast can be categorized into three groups on the basis of their salinity tolerance: a) Group-I that includes species like *Atriplex stocksii* (Khan & Rizvi, 1994), and *Zygophyllum simplex* (Khan & Ungar, 1997) which can germinate in up to 29.1 dS m^{-1} NaCl, b) Group-II that includes halophytes such as *Aeluropus lagopoides* (Gulzar & Khan, 2001), *Haloxylon stocksii* (Khan & Ungar, 1996), *Limonium stocksii* (Zia & Khan, 2002), and *Desmostachya bipinnata* (Gulzar *et al.* 2007) which have ability to tolerate up to 48.5 dS m^{-1} NaCl salinity at germination, and c) Group-III that includes species like *Arthrocnemum macrostachyum* (Khan & Gul, 1998), and *Cressa cretica* (Khan, 1999) which can germinate in up to 97 dS m^{-1} NaCl salinity. All test species appear to be moderately salt tolerant at germination and belong to Group-II, except *Suaeda fruticosa*, where no seed could germinate above 30 dS m^{-1} , hence can be included in Group-I.

Aeluropus lagopoides was the most tolerant, where comparatively more seeds germinated with higher seed germination rate at 50 dS m^{-1} in either salt type than other halophytic species. Gulzar & Khan (2001) reported identical results for *A. lagopoides*. Seeds of *Desmostachya bipinnata* and *Haloxylon stocksii* also germinated in up to 50 dS m^{-1} but rates as well as mean final germination in *D. bipinnata* were higher as compare to *H. stocksii*. *Suaeda fruticosa* was least salt tolerant among all species and showed no germination above 30 dS m^{-1} in both salts. Similar findings have been reported by Khan *et al.* (2006) for these species with seasalt salinity. However, Khan & Ungar (1998) in contrast reported that few seeds of *S. fruticosa* could germinate in up to 500 mM NaCl salinity. This difference might be attributed to genetic variability among populations. Therefore, a detailed study on intra-specific variation in seed germination responses among different populations of *S. fruticosa* is suggested.

There is little information is available on comparative influence of NaCl and seasalt on seed germination of halophytes (Joshi *et al.* 1995; Baskin & Baskin, 1998; Houle *et al.* 2001; Zia & Khan, 2002; Atia *et al.* 2006; Liu *et al.* 2006). Seed germination of *A. lagopoides* and *Desmostachya bipinnata* inhibited more in seasalt especially at higher concentration in comparison to *H. stocksii*. However, seed germination of *S. fruticosa* also reduced more in seasalt than NaCl especially above 10 dS m⁻¹. Zia & Khan (2002) also reported similar results for *Limonium stocksii*, where seasalt suppressed germination more as compared to NaCl. In contrast, Tirmizi *et al.* (1993) have reported that NaCl inhibits seed germination of *Hipophae rhamnoides* more in comparison to seasalt. Duan *et al.* (2003) also reported that NaCl inhibits germination of *Suaeda salsa* more than seasalt. However, De Villiers *et al.* (1994) and Houle *et al.* (2001) reported inhibitorier effect of seasalt than NaCl on seed germination. This can be attributed to the fact that seasalt is a combination of several salts with leading concentration of NaCl, and seasalt induced germination suppression might be a consequence of synergistic lowering of osmotic potential of the medium than NaCl alone. However, a thorough physiological investigation is needed to understand why seasalt is more inhibitory in some species and not in others.

In summary, halophytic vegetation of Karachi coast varied in their salinity tolerance range and shows various effect in response to NaCl and seasalt at germination stage. *Aeluropus lagopoides*, *Desmostachya bipinnata* (Monocots) and *Haloxylon stocksii* (Dicots) were found to be moderately salinity tolerant (50 dS m⁻¹), while *Suaeda fruticosa* (Dicots) was low salt-tolerance (30 dS m⁻¹) under both salt condition. Salinity tolerance of test species was in following order: *A. lagopoides* > *D. bipinnata* > *H. stocksii* > *S. fruticosa*. The effect of both salts (NaCl and Seasalt) on germination was also varied among species. Seasalt affected seed germination of *A. lagopoides*, *D. bipinnata* (Monocots) and *S. fruticosa* (Dicots) more adversely than NaCl especially at higher concentration. However, seed germination of *H. stocksii* (Dicots) was not significantly different in either salt type. Variation in rate of seed germination of *Desmostachya bipinnata* and *Suaeda fruticosa* under NaCl and seasalt present only at high doses of salinity, whereas difference was absent for *Aeluropus lagopoides* and *Haloxylon stocksii*.

References

- Adam, P. 1990. Salt marsh ecology. Cambridge university press, New York.
- Aronson, J.A. 1989. Haloph: A data base of salt tolerant plants of the world. Office of Arid Lands Studies. The University of Arizona, Tucson.
- Atia, A., K.B. Hamed, A. Debez and C. Abdely. 2006. Salt and seawater effects on the germination of *Crithmum maritimum*. In: Biosaline agriculture and salinity tolerance in plants. M. Ozturk, Y. Waisel, M.A. Khan, G. Gork. and B. Verlag. (Eds.) Switzerland. pp. 29-33.
- Baskin, J.M. and C.C. Baskin, 1998. The annual dormancy cycle in buried weed seeds: a continuum. *Biol. Sci.*, 35: 492-498.
- Benwahhoud, M., H. Jouad, M. Eddouks and B. Lyoussi. 2001. Hypoglycemic effect of *Suaeda fruticosa* in streptozotocin-induced diabetic rats. *J. Ethnopharmacol.*, 76: 35-38.
- Chapman, V.J. 1960. Salt marshes and salt deserts of the world. New York, New York: Interscience Publishers.
- Cope, T.A. 1982. Poaceae. In: *Flora of Pakistan*. E. Nasir and S.I. Ali (Eds.). University of Karachi: Karachi, Pakistan.

- DeVilliers, A.J., M.W. Van Rooyen, G.K. Theron, and H.A. Van De Venter. 1994. Germination of three Namaqualand pioneer species, as influenced by salinity, temperature and light. *Seed Sci. Technol.*, 22: 427-433.
- Duan, D., X. Liu, F. Feng, and C. Li. 2003. Effects of salinities on seed germination of halophytes *Suaeda salsa*. *Chin. Agric. Sci. Bull.*, 19: 168-172.
- Gulzar, S. and M.A. Khan, 2001. Seed germination of a halophytic grass *Aeluropus lagopoides*. *Ann. Bot.*, 87: 319-324.
- Houle, G., L. More, C.E. Reynolds and J.Siége. 2001. The effect of salinity on different developmental stages of an endemic annual plant *Aster laurentianus* (Asteraceae). *Am. J. Bot.*, 88: 62-67.
- Joshi, A.J. and A.S. Bhoite. 1988. Fluctuation of mineral ions in saline soils and halophytic grass *Aeluropus lagopoides* L. *Ann. Arid Zone*, 27: 191-196.
- Joshi, A.J., B.S. Mali, and H. Hingalajia. 1995. Halophytes A good source of forage production under salt stress conditions 1. *Salvadora persica*. In: Biology of Salt Tolerant Plants, M.A. Khan and I.A. Ungar (eds.). Department of Botany, University of Karachi. pp: 353-360.
- Khan, M.A. 1999. Comparative influence of salinity and temperature on the germination of subtropical halophytes. In: Halophyte Uses in different climates I: Ecological and Ecophysiological Studies. H. Lieth, M. Moschenko, M. Lohman, H.W. Koyro and A. Hamdy (eds.). Progress in Biometeorology, Backhuys Publishers, Leiden, 13: 77-88.
- Khan, M.A. and B. Gul. 1998. High salt tolerance in the germinating dimorphic seeds of *Arthrocnemum indicum*. *Int. J. Plant Sci.*, 159: 826-832.
- Khan, M.A. and B. Gul. 2002a. Some ecophysiological aspects of seed germination in halophytes. In: Halophyte Utilization and Regional Sustainable Development of Agriculture. X. Liu and M. Liu (eds.). Meteorological Press, Beijing. pp: 56-68.
- Khan, M.A. and B. Gul. 2002b. Salt tolerant plants of coastal sabkhas of Pakistan. In: Sabkha Ecosystems. H. Barth and B. Boer (eds.). Kluwer Academic Press, Netherlands.
- Khan, M.A. and B. Gul. 2006. Halophyte seed germination. In: Eco-physiology of High Salinity Tolerant Plants, M.A. Khan and D.J. Weber (eds.). Springer, pp: 11-30.
- Khan, M.A. and M. Qaiser. 2006. Halophytes of Pakistan: Distribution, Ecology, and Economic Importance. In: Sabkha Ecosystems. M.A. Khan, H. Barth, G.C. Kust and B. Boer (eds.). Vol. II: Springer, Netherlands, pp: 129-153.
- Khan, M.A. and Y. Rizvi. 1994. Effect of salinity, temperature and growth regulators on the germination and early seedling growth of *Atriplex griffithii* var. *stocksii*. *Can. J. Bot.*, 72: 475-479.
- Khan, M.A. and I.A. Ungar. 1984. The effect of salinity and temperature on germination of polymorphic seeds and growth of *Atriplex triangularis* Willd. *Am. J. Bot.*, 71: 481-489.
- Khan, M.A. and I.A. Ungar. 1996. Influence of salinity and temperature on the germination of *Haloxylon recurvum*. *Ann. Bot.*, 78: 547-551.
- Khan, M.A. and I.A. Ungar. 1997. Germination responses of the subtropical annual halophyte *Zygophyllum simplex*. *Seed Sci. Technol.*, 25: 83-91.
- Khan, M.A. and I.A. Ungar. 1998. Germination of salt tolerant shrub *Suaeda fruticosa* from Pakistan: Salinity and temperature responses. *Seed Sci. Technol.*, 26: 657-667.
- Khan, M.A., M.Z. Ahmed and A. Hameed. 2006. Effect of sea salt and L-ascorbic acid on the seed germination of halophytes. *J. Arid Environ.*, 65: 535-540.
- Khan, M.A., B. Gul and D.J. Weber. 2002. Effect of temperature, and salinity on the germination of *Sarcobatus vermiculatus*. *Biol. Plant.*, 45: 133-135.
- Liu, X., H. Qiao, W. Li, T. Tadano and M.A. Khan. 2006. Comparative effect of NaCl and Seawater on seed germination of *Suaeda salsa* and *Atriplex centralasiatica*. In: Biosaline agriculture and salinity tolerance in plants. M. Ozturk, Y. Waisel, M.A. Khan and G. Gork (eds.). Birkhauser Verlag, Switzerland, pp: 45-53.
- McMillon, C. 1988. Seed germination and seedling development of *Halophila decipiens*. Ostenfeld from Panama. *Aquat. Bot.*, 31: 169-176.

- Mooring, M.T., A.W. Cooper and E.D. Seneca. 1971. Seed germination response and evidence for height of ecophenes in *Spartina alterniflora* from North Carolina. *Am. J. Bot.*, 58: 48-56.
- Noe, G.B. and J.B. Zedler. 2000. Differential effects of four abiotic factors on the germination of salt marsh annuals. *Am. J. Bot.*, 87: 1679-1692.
- Rivers, W.G. and D.J. Weber. 1971. The influence of salinity and temperature on seed germination in *Salicornia bigelovii*. *Physiol. Plant.*, 24: 73-75.
- SPSS Inc. 2001. SPSS: SPSS 11.0 for Windows *Update*. SPSS Inc. USA.
- Stewart, R.R. 1972. An annotated catalogue of the vascular plants of West Pakistan and Kashmir. In: Flora of Pakistan, E, Nasir and S.I. Ali (Eds.). Department of Botany, University of Karachi. 1-1028.
- Tirmizi, S.A.S., K.M. Khan and S.A. Qadir. 1993. Study on salt tolerance of *Hippophae rhamnoides* L. during germination. *Pak. J. Sci. Ind. Res.*, 36: 252-257.
- Ungar, I.A. 1978. Halophyte seed germination. *Bot. Rev.*, 44: 233-264.
- Ungar, I.A. 1991. Ecophysiology of Vascular Halophytes. CRC Press, Boca Raton, F.L.
- Ungar, I.A. 1995. Seed germination and seed-bank ecology of halophytes. In: Seed development and germination, J. Kigel and G. Galili (eds.). Marcel and Dekker, New York. pp: 599-628.
- Yokoishi, T. and S. Tanimoto, 1994. Seed germination of the halophyte *Suaeda japonica* under salt stress. *J. Plant Res.*, 107: 385-388.
- Zia, S. and M.A. Khan. 2002. Comparative effect of NaCl and seawater on seed germination of *Limonium stocksii*. *Pak. J. Bot.*, 34: 345-350.

(Received for publication 18 September, 2006)

**PAKISTAN BOTANICAL SOCIETY
OFFICERS AND COUNCIL FOR 2006**

President

Dr. M. Ashraf
Department of Botany, University of Agriculture, Faisalabad.

Vice-President

Dr. S.M. Saqlain Naqvi
Department of Biochemistry, University of Arid Agriculture, Rawalpindi.

Dr. Muhammad Tahir Rajput
Department of Botany, University of Sindh, Jamshoro.

Dr. Mufakhira Jan Durrani
Department of Botany, University of Balochistan, Quetta.

Dr. Farrukh Hussain
Department of Botany, University of Peshawar, Peshawar.

Mr. Shafiq-ur-Rehman
Department of Botany, University of AJ&K Muzaffarabad.

Secretary/Treasurer

Dr. M. Yasin Ashraf
NIAB, Faisalabad.

Joint Secretary

Dr. Syeda Saleha Hassney
Department of Botany, University of Sindh, Jamshoro.

Chief Editor

Dr. Abdul Ghaffar,
Department of Botany, University of Karachi.

COUNCIL

The Officers of the Pakistan Botanical Society, the President and Secretary ex officio, Prof. Nuzhat Parveen, Shipowners College, Karachi; Dr. A. Qayyum Soomro, University of Sindh, Jamshoro; Dr. Shahida Hasnain, University of the Punjab, Lahore; Dr. Manzoor H. Soomro, PSF, Islamabad; Dr. Fazle Malik Sarim, University of Peshawar; Dr. A. Razzak Mahar, Shah Abdul Latif University, Khairpur; Dr. Tayyab Hussain Bukhari, Govt. College, Toba Tek Singh.

	Page
Muhammad Ashraf and Habib-ur-Rehman Athar - Preface	1357
Sayed Anwar-ul-Haque - Salinity problems and crop production in coastal regions of Bangladesh - Review Article	1359
Abdallah Atia, Ahmed Debez, Mokded Rabhi, Habib-ur-Rehman Athar, and Chedly Abdelly - Alleviation of salt-induced seed dormancy in the perennial halophyte <i>Crithmum maritimum</i> L. (<i>Apiaceae</i>)	1367
Shahid Umar - Alleviating adverse effects of water stress on yield of sorghum, mustard and groundnut by potassium application	1373
M.U. Shirazi, M.A. Khan, Mukhtiar Ali, S. M. Mujtaba, S. Mumtaz, Muhammad Ali, B. Khanzada, M.A. Halo, M. Rafique, J. A. Shah, K.A. Jafri, and N. Depar - Growth performance and nutrient contents of some salt tolerant multipurpose tree species growing under saline environment	1381
Mumtaz Hussain, Muhammad Sajid Aqeel Ahmad and Abida Kausar - Effect of lead and chromium on growth, photosynthetic pigments and yield components in mash bean [<i>Vigna mungo</i> (L.) Hepper]	1389
Bechir Ben-Rouina, Chedlia Ben Ahmed, Habib-ur-Rehman Athar, and M. Boukhriss - Water relations, proline accumulation and photosynthetic activity in olive tree (<i>Olea europaea</i> L. cv "Chemlali") in response to salt stress	1397
Ameer Khan, Muhammad Sajid Aqeel Ahmad, Habib-ur-Rehman Athar and Muhammad Ashraf - Interactive effect of foliar applied ascorbic acid and salt stress on wheat (<i>Triticum aestivum</i> L.) at seedling stage	1407
A. R. Gurmani, A. Bano and Muhammad Salim - Effect of growth regulators on growth, yield and ions accumulation of rice (<i>Oryza sativa</i> L.) under salt stress	1415
Muhammad Asif, Mehboob-ur-Rahman and Yusuf Zafar - Genotyping analysis of six maize (<i>Zea mays</i> L.) hybrids using DNA fingerprinting technology	1425
Sezgin Celik - Studies on the germination of three endangered <i>Centurea</i> species	1431
Vishandaz, Zia-ul-Hassan, Muhammad Arshad and Ahmad Naqi Shah - Phosphorus fertigation at first irrigation due to its unavailability at sowing time prevents yield losses in <i>Triticum aestivum</i> L.	1439
M. Kashif Shahzad Sarwar, Ihsan Ullah, Mehboob-ur-Rahman, M. Yasin Ashraf and Yusuf Zafar - Glycinebetaine accumulation and its relation to yield and yield components in cotton genotypes grown under water deficit condition	1449
Muhammad Sajid Aqeel Ahmad, Qasim Ali, Rohina Bashir, Farrukh Javed And Ambreen Khadija Alvi - Time course changes in ionic composition and total soluble carbohydrates in two barley cultivars at seedling stage under salt stress	1457
Tayyaba Shaheen, Mehboob-ur-Rahman and Yusuf Zafar - Chloroplast rps8 gene of cotton reveals the conserved nature through out plant taxa	1467
Chedlia Ben-Ahmed, B. Ben-Rouina, Habib-ur-Rehman Athar, M. Boukhriss - Olive tree (<i>Olea europaea</i> L. cv. "Chemlali") under salt stress: water relations and ions content	1477
Jim Morris, John Collopy and Khalid Mahmood - Canopy conductance and water use in <i>Eucalyptus</i> plantations	1485
B. Shaharona, Riffat Bibi, Muhammad Arshad, Zahir Ahmed Zahir, and Zia-ul-Hassan - 1-Aminocyclopropane-1-carboxylate (ACC)-deaminase rhizobacteria extenuates ACC-induced classical triple response in etiolated pea seedlings	1491
Roubina Kausar, Habib-ur-Rehman Athar, and Muhammad Ashraf - Chlorophyll fluorescence can be used as a potential indicator for rapid assessment of water stress tolerance in canola (<i>Brassica napus</i> L.)	1501

Uzma Farooq and Asghari Bano - Effect of abscisic acid and chlorocholine chloride on nodulation and biochemical content of <i>Vigna radiata</i> L. under water stress	1511
Ruya Yilmaz, Serdal Sakcali, Celal Yarci, Ahmet Aksoy and Munir Ozturk - Use of <i>Aesculus hippocastanum</i> L. as a biomonitor of heavy metal pollution.	1519
Tariq Aziz, Rahmatullah, Muhammad Aamer Maqsood, Mukkram. A. Tahir, Iftikhar Ahmad and Mumtaz A. Cheema - Phosphorus utilization by six Brassica cultivars (<i>Brassica juncea</i> L.) from tri-calcium phosphate; a relatively insoluble P compound	1529
Gul Sanat Shah Khattak, Iqbal Saeed And Tila Muhammad - Breeding for heat tolerance in mungbean (<i>Vigna radiata</i> (L.) Wilczek)	1539
Nazila Azhar, M. Yasin Ashraf, M. Hussain and F. Hussain - Phytoextraction of lead (Pb) by EDTA application through sunflower (<i>Helianthus annuus</i> L.) cultivation: Seedling growth studies.	1551
Muhammad Ibrahim, Ambreen Anjum, Nabeela Khaliq, Muhammad Iqbal and Habib-ur-Rehman Athar - Four foliar applications of glycinebetaine did not alleviate adverse effects of salt stress on growth of sunflower	1561
Muhammad Aslam, Iftikhar A. Khan, Muhammad Saleem and Zulfiqar Ali - Assessment of water stress tolerance in different maize accessions at germination and at early growth stage	1571
Nudrat Aisha Akram, Muhammad Shahbaz, Habib-ur-Rehman Athar, and Muhammad Ashraf - Morpho-physiological responses of two differently adapted populations of <i>Cynodon dactylon</i> (L.) Pers. and <i>Cenchrus ciliaris</i> L. to salt stress	1581
M. Yasin Ashraf, Kalsoom Akhter, F. Hussain and Javed Iqbal - Screening of different accessions of three potential grass species from Cholistan desert for salt tolerance	1589
Nabila Tabbasam, Mehboob-ur-Rahman and Yusuf Zafar - DNA-based genotyping of sorghum hybrids	1599
Abdul Hameed, Muhammad Zaheer Ahmed and Muhammad Ajmal Khan - Comparative effects of NaCl and seasalt on seed germination of coastal halophytes	1605
Tanwir Ahmad Malik, Sana-Ullah, and Samina Malik - Genetic linkage studies of drought tolerant and agronomic traits in cotton	1613
Qasim Ali, Habib-ur-Rehman Athar and Muhammad Ashraf - Influence of exogenously applied brassinosteroids on the mineral nutrient status of two wheat cultivars grown under saline conditions	1621
M. A. Khan, M.U. Shirazi, Mukhtiar Ali, S. Mumtaz, A. Sherin, and M. Y. Ashraf - Comparative performance of some wheat genotypes growing under saline water	1633
Naeem Iqbal and Muhammad Yasin Ashraf - Does seed treatment with glycinebetaine improve germination rate and seedling growth of sunflower (<i>Helianthus annuus</i> L.) under osmotic stress	1641
Irfan Afzal, Shahzad Maqsood Ahmad Basra, Amjad Hameed and Muhammad Farooq - Physiological enhancements for alleviation of salt stress in wheat	1649
Muhammad Saeed Akram, Qasim Ali, Habib-ur-Rehman Athar and Ahmed Saeed Bhatti - Ion uptake and distribution in <i>Panicum antidotale</i> Retz. under salt stress	1661
Sulman Shafeeq, Mehboob-ur-Rahman and Yusuf Zafar - Genetic variability of different wheat (<i>Triticum aestivum</i> L.) genotypes/cultivars under induced water stress	1671

Cont'd. on page 1770

Ihsan Ullah, Mehboob-ur-Rahman and Yusuf Zafar - Genotypic variation for drought tolerance in cotton (<i>Gossypium hirsutum</i> L.): seed cotton yield responses	1679
Ismat Nawaz, Zia-Ul-Hassan, Atta Muhammad Ranjha and Muhammad Arshad - Exploiting genotypic variation among fifteen maize genotypes of Pakistan for potassium uptake and use efficiency in solution culture	1689
Munir Ahmad, Zahid Akram, Muhammad Munir and Muhammad Rauf - Physio-morphic response of wheat genotypes under rainfed conditions	1697
Qasim Ali, Habib-ur-Rehman Athar and Muhammad Ashraf - Ion transport in four canola cultivars as influenced by salt stress	1703
S.F. Afzali, M.A. Hajabbasi, H. Shariatmadari, K. Razmjoo, and A.H. Khoshgofarmanesh - Comparative adverse effects of PEG- or NaCl-induced osmotic stress on germination and early seedling growth of a potential medicinal plant <i>Matricaria chamomilla</i>	1709
Mukkram Ali Tahir, Rahmatullah, Tariq Aziz, M.Ashraf, Shamsa Kanwal and Muhammad Aamer Maqsood - Beneficial effects of silicon in wheat (<i>Triticum aestivum</i> L.) under salinity stress	1715
Shamyla Nawazish, Mansoor Hameed and Shaista Naurin - Leaf anatomical adaptations of <i>Cenchrus ciliaris</i> L. from the salt range, Pakistan against drought stress	1723
Farah Naz, Shafqat Farooq, Rubina Arshad, Muhammad Afzaal and Muhammad Akram - Discriminating upland and lowland rice genotypes through proteomic approach	1731
Muhammad Akram, Shafqat Farooq, Muhammad Afzaal, Farah Naz, and Rubina Arshad - Chlorophyll fluorescence in different wheat genotypes grown under salt stress	1739
Muhammad Afzaal, Shafqat Farooq, Muhammad Akram, Farah. Naz, Rubina Arshad and Asghari Bano - Differences in agronomic and physiological performance of various wheat genotypes grown under saline conditions	1745
Rubina Arshad, Shafqat Farooq and Farooq-E-Azam - Rhizospheric bacterial diversity: Is it partly responsible for water deficiency tolerance in wheat?	1751
Shamaila Rafiq, Tahira Iqbal, Amjad Hameed, Zulfiqar Ali Rafiqi and Naila Rafiq - Morphobiochemical analysis of salinity stress response of wheat	1759

Published by the Pakistan Botanical Society.

Printed at Karachi University Press, Karachi-75270, Pakistan. Phone: 9243186, 4969237.

Copies available from the Chief Editor, Pakistan Journal of Botany, Department of Botany, University of Karachi, Karachi-75270, Pakistan. Phone: 4387867.

Price: Each issue Rs. 100/ US\$20; Vol. Rs. 600/US\$120 or in exchange.